

COMPLETENESS ON THE WORM DOMAIN AND THE MÜNTZ–SZÁSZ PROBLEM FOR THE BERGMAN SPACE

STEVEN G. KRANTZ, MARCO M. PELOSO, AND CATERINA STOPPATO

ABSTRACT. In this paper we are concerned with the problem of completeness in the Bergman space of the worm domain \mathcal{W}_μ and its truncated version \mathcal{W}'_μ . We determine some orthogonal systems and show that they are not complete, while showing that the union of two particular of such systems is complete.

In order to prove our completeness result we introduce the *Müntz–Szász problem* for the 1-dimensional Bergman space of the disk $\{\zeta : |\zeta - 1| < 1\}$ and find a sufficient condition for its solution.

INTRODUCTION

The Diederich–Fornaess worm domain was introduced in [DF77] and is defined for a given $\mu > 0$ as

$$\mathcal{W}_\mu = \{(z_1, z_2) \in \mathbb{C}^2 : |z_1 - e^{i \log |z_2|^2}| < 1 - \varphi(\log |z_2|^2)\}, \quad (1)$$

where $\varphi : [-A, A] \rightarrow [0, 1]$ is a smooth, convex, even function that vanishes identically in $[-\mu, \mu]$, with $A > \mu$, $\varphi(A) = 1$, and increasing on $[\mu, A]$. As a result, \mathcal{W}_μ is smooth, pseudoconvex and strictly pseudoconvex at all points $(z_1, z_2) \in \partial \mathcal{W}_\mu$ with $z_1 \neq 0$. See [CS01] for a thorough discussion of basic properties of the worm. The worm turned out to be of fundamental importance in the theory of geometric analysis in several complex variables, see [Kis91], [Bar92], [Chr96], [KP08a], [KP08b], [KPS15] and references therein.

For computational purposes, \mathcal{W}_μ is often truncated to the non-smooth bounded domain

$$\mathcal{W}'_\mu = \{(z_1, z_2) \in \mathbb{C}^2 : |z_1 - e^{i \log |z_2|^2}| < 1, \quad |\log |z_2|^2| < \mu\}, \quad (2)$$

that is, one replaces the function φ with the characteristic function of the complement of the interval $[-\mu, \mu]$.

In the discussion that follows, we let Ω denote either domain \mathcal{W}_μ or \mathcal{W}'_μ .

In this study we are concerned with the question of finding orthogonal and complete sets in $A^2(\Omega)$. As is well known, when $\mu \geq \pi$, Ω has non-trivial *Nebenhülle*. Here the *Nebenhülle* is understood to be the interior of the connected component that contains $\overline{\Omega}$ of the intersection of all domains of holomorphy containing $\overline{\Omega}$. We first show that this easily implies that the closure in $A^2(\Omega)$ of the holomorphic polynomials is a proper subset of $A^2(\Omega)$.

2010 *Mathematics Subject Classification.* 32A25, 32A36, 30H20.

Key words and phrases. Bergman kernel, Bergman projection, worm domain.

Second author supported in part by the 2010-11 PRIN grant *Real and Complex Manifolds: Geometry, Topology and Harmonic Analysis* of the Italian Ministry of Education (MIUR).

Third author supported by the same PRIN grant; by the FIRB grant *Differential Geometry and Geometric Function Theory* of the MIUR; and by the research group GNSAGA of the Istituto Nazionale di Alta Matematica.

Thus we are led to consider sets of suitable “monomials” in Ω that can be defined as the holomorphic continuation of non-integral powers z_1^η , when $z = (z_1, z_2)$ is initially restricted to $\Delta \times \{z_2 : |z_2| = 1\}$, where we denote by Δ the disk $\{\zeta \in \mathbb{C} : |\zeta - 1| < 1\}$.

We determine some orthogonal sets $\{H_{2k,j}, k, j \in \mathbb{Z}, k \geq 0\}$, and $\{H_{2k+1,j}, m, j \in \mathbb{Z}, k \geq 0\}$ (see Corollary 1.5), and show that their union determines a complete set in $A^2(\mathcal{W}'_\mu)$ when $\mu > 0$ (Theorem 3.1). We also show that each of the two systems, however, is not complete (Proposition 3.2).

In order to prove our completeness result, Theorem 3.1, we prove a result of independent interest, Theorem 2.1. We naturally use the name *the Müntz–Szász problem for the Bergman space* for the question of characterizing the sequences $\{\lambda_j\}$ in the right half-plane for which the sets of powers $\{\zeta^{\lambda_j-1}\}$ form a complete set in $A^2(\Delta)$.

The classical Müntz–Szász theorem deals with the completeness of sets of powers $\{t^{\lambda_j-\frac{1}{2}}\}$ in $L^2([0, 1])$, where again λ_j is in the right half plane. The solution was provided by C. Müntz [Mün14] and by O. Szász [Sz16] in two separate papers, where they showed that the set $\{t^{\lambda_j-\frac{1}{2}}\}$ is complete $L^2([0, 1])$ if and only if $\sum_{j=1}^{+\infty} (1 + |\lambda_j|^2)^{-1} \operatorname{Re} \lambda_j = \infty$ (see also [PW87] or [Rud66] for a more accessible reference).

We find a sufficient condition for the solution of the Müntz–Szász problem for the Bergman space $A^2(\Delta)$ and use it to prove our completeness result for $A^2(\mathcal{W}'_\mu)$.

Finally we show that the complete set $\{H_{k,j}, k, j \in \mathbb{Z}, k \geq 0\}$, is not a basis (Theorem 3.3).

1. ORTHOGONAL SETS IN $A^2(\mathcal{W}'_\mu)$

Let $\mu > 0$ and consider the domain \mathcal{W}'_μ . The problem we address here is to find a, possibly complete, orthonormal system for $A^2(\mathcal{W}'_\mu)$ and consequently have a way to obtain an expression for the Bergman kernel.

In the analysis on the worm domains \mathcal{W}_μ and \mathcal{W}'_μ a special role is played by the functions

$$E_\eta(z) = e^{\eta L(z)}, \quad (3)$$

where

$$L(z) = \log(z_1 e^{-i \log |z_2|^2}) + i \log |z_2|^2, \quad (4)$$

and \log denotes the principal branch of the logarithm, so that

$$E_\eta(z_1, z_2) = (z_1 e^{-i \log |z_2|^2})^\eta e^{i \eta \log |z_2|^2}.$$

The function L is well defined and holomorphic in a domain containing $\cup_\mu \mathcal{W}'_\mu$ (see [KPS15], Lemma 1.2 and Proposition 1.3). Moreover, we point out that the fiber of \mathcal{W}'_μ over each $z_1 \in D(0, 2) \setminus \{0\}$ is not connected and that $L(z)$ is locally constant in z_2 , but not constant. The same happens with $E_\eta(z)$ for $\eta \in \mathbb{C} \setminus \mathbb{Z}$, while $E_k(z) = z_1^k$ for all $k \in \mathbb{Z}$, $z \in \mathcal{W}'_\mu$. Hence the functions E_η are the analytic continuation to \mathcal{W}'_μ of the monomial z_1^η defined in $\mathcal{W}'_{\pi/2}$ using the principal branch of the logarithm.

It is well known that the functions that are holomorphic in a neighborhood of the closure $\overline{\mathcal{W}_\mu}$ are not dense in $A^2(\mathcal{W}_\mu)$. Since a proof of this fact does not explicitly appear in the literature, we prove the following result that applies to both domains \mathcal{W}_μ and \mathcal{W}'_μ .

Proposition 1.1. *Let $\mu \geq 2\pi$ and let $A^2(\overline{\mathcal{W}_\mu})$ denote the closure in $A^2(\mathcal{W}_\mu)$ of the functions that are holomorphic in a neighborhood of \mathcal{W}_μ . Then, if $f \in A^2(\overline{\mathcal{W}_\mu})$, then f is holomorphic on $\widehat{\mathcal{W}_\mu}$, where*

$$\widehat{\mathcal{W}_\mu} \supseteq \bigcup_{-\mu \leq a \leq \mu - 2\pi} \{(z_1, z_2) : a < \log |z_2|^2 < a + 2\pi, |z_1 - e^{ia}| < 1\}. \quad (5)$$

Therefore $A^2(\overline{\mathcal{W}_\mu}) \subsetneq A^2(\mathcal{W}_\mu)$.

The same conclusions hold true with \mathcal{W}'_μ in place of \mathcal{W}_μ .

In particular, the polynomials are not dense in either $A^2(\mathcal{W}_\mu)$ or $A^2(\mathcal{W}'_\mu)$. By contrast, D. Catlin [Cat80] showed that for every smoothly bounded pseudoconvex domain Ω , the holomorphic functions in $C^\infty(\overline{\Omega})$ are dense in $A^2(\Omega)$.

Proof. It suffices to prove the result in the case of \mathcal{W}'_μ , since the same argument can be repeated *verbatim* for \mathcal{W}_μ .

Suppose f is holomorphic in a neighborhood of $\overline{\mathcal{W}'_\mu}$ and let

$$\mathcal{E}_a = \{(0, z_2) : a \leq \log |z_2|^2 \leq a + 2\pi\} \bigcup \{(z_1, z_2) : \log |z_2|^2 = a \text{ or } a + 2\pi, |z_1 - e^{ia}| \leq 1\}.$$

Set

$$F_a(z_1, z_2) = \frac{1}{2\pi i} \int_\gamma \frac{f(z_1, \zeta)}{\zeta - z_2} d\zeta,$$

where γ is the oriented boundary of the annulus $\{z_2 \in \mathbb{C} : a < \log |z_2|^2 < a + 2\pi\}$. Then F_a is holomorphic on the set

$$\tilde{\mathcal{E}}_a = \{(z_1, z_2) : a < \log |z_2|^2 < a + 2\pi, |z_1 - e^{ia}| < 1\}.$$

However, $F_a(0, z_2) = f(0, z_2)$, since f is holomorphic in a neighborhood of $\{(0, z_2) : |\log |z_2|^2| \leq \mu\}$ and $-\mu \leq a \leq \mu - 2\pi$ implies $a + 2\pi \leq \mu$. It follows that F_a is a holomorphic extension of f to the set $\tilde{\mathcal{E}}_a$ and thus f extends holomorphically to an open set $\widehat{\mathcal{W}'_\mu}$ containing the right-hand side of (5).

Now suppose that $\{f_n\}$ are holomorphic in a neighborhood of $\overline{\mathcal{W}'_\mu}$ and that $f_n \rightarrow f$ in $A^2(\mathcal{W}'_\mu)$. Then $f_n \rightarrow f$ uniformly on the compact sets $\{\log |z_2|^2 = a, a + 2\pi; |z_1 - e^{ia}| \leq 1 - \delta\}$, for $-\mu < a < \mu - 2\pi$. By Cauchy's formula, $\{f_n\}$ is Cauchy in uniform norm also on the sets

$$\{a \leq \log |z_2|^2 \leq a + 2\pi, |z_1 - e^{ia}| \leq 1 - \delta\}.$$

Therefore f extends holomorphically to the set on the right-hand side of (5).

Finally, the functions E_η with η not an integer cannot be extended holomorphically to any of the sets $\tilde{\mathcal{E}}_a$, so that $A^2(\overline{\mathcal{W}'_\mu}) \subsetneq A^2(\mathcal{W}'_\mu)$.

It is immediate to check that the arguments above apply to the case of \mathcal{W}_μ as well. \square

Thus, we are led to consider the set of “monomials” of the form $\{E_{\eta_j}(z)z_2^j\}_{j \in \mathbb{Z}}$ and ask whether these are orthogonal, and/or complete, for some choice of values $\eta_j \in \mathbb{C}$.

We denote by dA the Lebesgue measure in the complex plane.

Lemma 1.2. *Let $\operatorname{Re} \alpha, \operatorname{Re} \beta > -1$. Then*

$$\int_\Delta \zeta^\alpha \overline{\zeta}^\beta dA(\zeta) = \pi \frac{\Gamma(\alpha + \overline{\beta} + 2)}{\Gamma(\alpha + 2)\Gamma(\overline{\beta} + 2)}.$$

In particular, ζ^α and ζ^β are never orthogonal to each other in $A^2(\Delta)$.

Proof. We have

$$\begin{aligned}
\int_{\Delta} \zeta^\alpha \overline{\zeta^\beta} dA(\zeta) &= \int_0^2 \int_{-\cos^{-1}(r/2)}^{\cos^{-1}(r/2)} (re^{i\theta})^\alpha (re^{-i\theta})^{\overline{\beta}} d\theta r dr \\
&= \int_0^2 r^{\alpha+\overline{\beta}+1} \int_{-\cos^{-1}(r/2)}^{\cos^{-1}(r/2)} e^{i\theta(\alpha-\overline{\beta})} d\theta dr \\
&= \frac{2}{i(\alpha-\overline{\beta})} \int_0^2 r^{\alpha+\overline{\beta}+1} \sinh(\cos^{-1}(r/2)i(\alpha-\overline{\beta})) dr \\
&= \frac{4}{i(\alpha-\overline{\beta})} \int_0^{\pi/2} (2\cos s)^{\alpha+\overline{\beta}+1} \sin s \sinh(is(\alpha-\overline{\beta})) ds \\
&= \frac{2^{\alpha+\overline{\beta}+3}}{\alpha+\overline{\beta}+2} \int_0^{\pi/2} (\cos s)^{\alpha+\overline{\beta}+2} \cosh(is(\alpha-\overline{\beta})) ds \\
&= \frac{2^{\alpha+\overline{\beta}+3}}{\alpha+\overline{\beta}+2} \int_0^{\pi/2} (\cos s)^{\alpha+\overline{\beta}+2} \cos(s(\alpha-\overline{\beta})) ds.
\end{aligned}$$

Now we use [GR96, (9) p. 391] and, denoting by B the beta function, we obtain that

$$\begin{aligned}
\int_{\Delta} \zeta^\alpha \overline{\zeta^\beta} dA(\zeta) &= \frac{\pi}{(\alpha+\overline{\beta}+2)(\alpha+\overline{\beta}+3)} \cdot \frac{1}{B(\alpha+2, \overline{\beta}+2)} \\
&= \pi \frac{\Gamma(\alpha+\overline{\beta}+2)}{\Gamma(\alpha+2)\Gamma(\overline{\beta}+2)},
\end{aligned}$$

as we wished to prove. \square

For a given bounded domain Ω in \mathbb{C}^2 that is rotationally invariant¹ in the second variable z_2 , such as \mathcal{W}_μ and \mathcal{W}'_μ , using the Fourier expansion in z_2 , the Bergman space $A^2(\Omega)$ decomposes as an orthogonal sum

$$A^2(\Omega) = \bigoplus_{j \in \mathbb{Z}} \mathcal{H}^j. \quad (6)$$

Here

$$\mathcal{H}^j = \{F \in A^2(\Omega) : F(z_1, z_2) = f(z_1, |z_2|)z_2^j\},$$

where f is holomorphic in z_1 and locally constant in $|z_2|$. The orthogonal projection of A^2 onto \mathcal{H}^j is given by

$$Q_j F(z_1, z_2) = \frac{1}{2\pi} \int_{-\pi}^{\pi} F(z_1, e^{it}z_2) e^{-ijt} dt.$$

Then we set

$$f(z_1, |z_2|) = \frac{Q_j F(z_1, z_2)}{z_2^j},$$

and observe that the right-hand side is holomorphic in Ω , but depends only on the modulus of z_2 . Hence f is locally constant in $|z_2|$. In general, it will be constant only if the fibers over a point (z_1, z_2) in Ω with z_1 fixed, is a connected set in the z_2 -plane.

¹These are called *Hartogs* domains.

Let us go back to the case of \mathcal{W}'_μ and let $F, G \in \mathcal{H}^j$, $F(z_1, z_2) = f(z_1, |z_2|)z_2^j$ and $G(z_1, z_2) = g(z_1, |z_2|)z_2^j$. We have

$$\begin{aligned} \langle F, G \rangle_{A^2(\mathcal{W}'_\mu)} &= \int_{|\log |z_2|^2| < \mu} \int_{|z_1 - e^{i \log |z_2|^2}| < 1} f(z_1, |z_2|) \overline{g(z_1, |z_2|)} |z_2|^{2j} dA(z_1) dA(z_2) \\ &= 2\pi \int_{|\log r^2| < \mu} \int_{|z_1 - e^{i \log r^2}| < 1} f(z_1, r) \overline{g(z_1, r)} dA(z_1) r^{2j+1} dr \\ &= \pi \int_{|s| < \mu} \int_{|z_1 - e^{is}| < 1} f(z_1, e^{s/2}) \overline{g(z_1, e^{s/2})} dA(z_1) e^{s(j+1)} ds \\ &= \pi \int_{\Delta} \int_{|s| < \mu} f(\zeta e^{is}, e^{s/2}) \overline{g(\zeta e^{is}, e^{s/2})} e^{s(j+1)} ds dA(\zeta). \end{aligned} \quad (7)$$

Let

$$\nu = \pi/2\mu. \quad (8)$$

be the reciprocal of the winding number of \mathcal{W}'_μ , as also defined in [Bar92] and let h be the entire function

$$h(z) = \frac{\sinh[\mu(j+1+iz)]}{j+1+iz}.$$

Define

$$\gamma_{\alpha\beta} = h(\alpha - \overline{\beta}). \quad (9)$$

Proposition 1.3. *Let $\mu > 0$. For $\alpha \in \mathbb{C}$ and $j \in \mathbb{Z}$ let $F_{\alpha,j}(z_1, z_2) = E_\alpha(z)z_2^j$. Then $F_{\alpha,j} \in A^2(\mathcal{W}'_\mu)$ if and only if $\operatorname{Re} \alpha > -1$. Moreover, if $\operatorname{Re} \alpha, \operatorname{Re} \beta > -1$ then*

$$\langle F_{\alpha,j}, F_{\beta,j} \rangle_{A^2(\mathcal{W}'_\mu)} = (2\pi)^2 \gamma_{\alpha\beta} \frac{\Gamma(\alpha + \overline{\beta} + 2)}{\Gamma(\alpha + 2)\Gamma(\overline{\beta} + 2)}.$$

In particular, $\langle F_{\alpha,j}, F_{\beta,j} \rangle_{A^2(\mathcal{W}'_\mu)} = 0$ if and only if

$$\alpha - \overline{\beta} = 2k\nu + i(j+1) \quad \text{with } k \in \mathbb{Z} \setminus \{0\}. \quad (10)$$

Proof. We compute $\langle F_{\alpha,j}, F_{\beta,j} \rangle_{A^2(\mathcal{W}'_\mu)}$. Starting from (7) we obtain

$$\begin{aligned} \langle F_{\alpha,j}, F_{\beta,j} \rangle_{A^2(\mathcal{W}'_\mu)} &= \pi \int_{\Delta} \int_{|s| < \mu} E_\alpha(\zeta e^{is}, e^{s/2}) \overline{E_\beta(\zeta e^{is}, e^{s/2})} e^{s(j+1)} ds dA(\zeta) \\ &= \pi \int_{\Delta} \zeta^\alpha \overline{\zeta}^\beta \int_{|s| < \mu} e^{is(\alpha - \overline{\beta})} e^{s(j+1)} ds dA(\zeta) \\ &= 2\pi \gamma_{\alpha\beta} \int_{\Delta} \zeta^\alpha \overline{\zeta}^\beta dA(\zeta), \end{aligned} \quad (11)$$

where

$$\begin{aligned} \gamma_{\alpha\beta} &:= \frac{1}{2} \int_{|s| < \mu} e^{s(j+1+i(\alpha - \overline{\beta}))} ds \\ &= \begin{cases} \frac{\sinh(\mu(j+1+i(\alpha - \overline{\beta})))}{j+1+i(\alpha - \overline{\beta})} & \text{if } j+1+i(\alpha - \overline{\beta}) \neq 0 \\ \mu & \text{if } j+1+i(\alpha - \overline{\beta}) = 0, \end{cases} \end{aligned}$$

as claimed.

Therefore, $\gamma_{\alpha\beta} = 0$ if and only if

$$\mu(j+1+i(\alpha-\bar{\beta})) = k\pi i \quad \text{for } k \in \mathbb{Z} \setminus \{0\},$$

that is,

$$\alpha - \bar{\beta} = 2k\nu + i(j+1) \quad \text{for } k \in \mathbb{Z} \setminus \{0\}. \quad (12)$$

Notice that, when $\alpha = \beta$, the previous computation gives

$$\begin{aligned} \|E_\alpha(z)z_2^j\|_{A^2(\mathcal{W}'_\mu)}^2 &= 2\pi\gamma_{\alpha,\alpha} \int_{\Delta} |\zeta^\alpha|^2 dA(\zeta) \\ &= 2\pi\gamma_{\alpha,\alpha} \int_{|\zeta|<1} e^{2[\operatorname{Re} \alpha \log |\zeta-1| - \operatorname{Im} \alpha \arg(\zeta-1)]} dA(\zeta) \\ &= 2\pi\gamma_{\alpha,\alpha} \int_{|\zeta|<1} |\zeta-1|^{2\operatorname{Re} \alpha} e^{-2\operatorname{Im} \alpha \arg(\zeta-1)} dA(\zeta), \end{aligned}$$

which is finite if and only if $\operatorname{Re} \alpha > -1$. This proves the first part of the statement. The second part now follows from Lemma 1.2. \square

The following corollaries now follow at once.

Corollary 1.4. *Then, for $\operatorname{Re} \alpha > -1$, we have that $F_{\alpha,j} \in A^2(\mathcal{W}'_\mu)$ and*

$$\|F_{\alpha,j}\|_{A^2(\mathcal{W}'_\mu)}^2 = (2\pi)^2 \frac{\sinh[\mu(j+1-2\operatorname{Im} \alpha)]}{j+1-2\operatorname{Im} \alpha} \frac{\Gamma(2+2\operatorname{Re} \alpha)}{|\Gamma(2+\alpha)|^2}.$$

For $c_0 > -1$, and $\ell = 0, 1, 2, \dots$ we set

$$H_{\ell,j}(z_1, z_2) = E_{c_0+\nu\ell+i(j+1)/2}(z)z_2^j. \quad (13)$$

Corollary 1.5. *For $\mu > 0$, each of the two sets*

$$\{H_{2k,j}, j \in \mathbb{Z}, k = 0, 1, 2, \dots\}, \quad \text{and} \quad \{H_{2k+1,j}, j \in \mathbb{Z}, k = 0, 1, 2, \dots\}, \quad (14)$$

is an orthogonal system in $A^2(\mathcal{W}'_\mu)$.

2. THE MÜNTZ–SZÁSZ PROBLEM FOR THE BERGMAN SPACE

In endeavoring to establish whether the system $\{F_{\alpha,j}\}$ is complete we are led to consider the Müntz–Szász problem for the Bergman space.

Recall that we set $\Delta = \{\zeta : |\zeta-1| < 1\}$. We consider a set of functions $\{\zeta^{\lambda_k}\}$, $k = 1, 2, \dots$ and would like to find a necessary and sufficient condition for this set to be a *complete set* in $A^2(\Delta)$, that is, its linear span to be dense in $A^2(\Delta)$.

Theorem 2.1. *Let \mathcal{S} be the subset of $A^2(\Delta)$ whose elements are the functions ζ^{λ_k} for $k = 0, 1, 2, \dots$, where $\lambda_k = ak + c_0 + ib$, $0 < a < 1$, $c_0 > -1$ and $b \in \mathbf{R}$. Then \mathcal{S} is a complete set in $A^2(\Delta)$.*

Proof. Consider the biholomorphic map $C : \mathcal{U} \rightarrow \Delta$ given by

$$C(\omega) = \frac{2i}{i + \omega}$$

of the upper half plane \mathcal{U} onto Δ . Then

$$T : A^2(\Delta) \ni f \mapsto (f \circ C)C' \in A^2(\mathcal{U})$$

is a surjective isometry. Next, by the Paley–Wiener theorem, the Fourier transform \mathcal{F} , given by

$$(\mathcal{F}g)(\xi) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} g(t)e^{-i\xi t} dt,$$

provides a surjective isometry of $A^2(\mathcal{U})$ onto $L^2((0, +\infty), d\xi/\xi)$, see e.g. [BBG⁺04]. Therefore, $\{\zeta^{\lambda_k}\}$ will be complete in $A^2(\Delta)$ if and only if $\{\mathcal{F}(T\zeta^{\lambda_k})\}$ is complete in $L^2((0, +\infty), d\xi/\xi)$.

Now,

$$T(\zeta^\lambda)(\omega) = -\frac{(2i)^{\lambda+1}}{(i + \omega)^{\lambda+2}},$$

while

$$\begin{aligned} \mathcal{F}\left((u+i)^{-(\lambda+2)}\right)(\xi) &= \frac{1}{2\pi} \int_{-\infty}^{+\infty} \frac{1}{(u+i)^{\lambda+2}} e^{-i\xi u} du \\ &= \frac{1}{i^{\lambda+2}\Gamma(\lambda+2)} \xi^{\lambda+1} e^{-\xi} \chi_{(0,+\infty)}(\xi), \end{aligned}$$

(see also [DGGMR07], Lemma 1).

Therefore,

$$\mathcal{F}(T(\zeta^\lambda))(\xi) = -\frac{(2\xi)^{\lambda+1}e^{-\xi}}{i\Gamma(\lambda+2)} \chi_{(0,+\infty)}(\xi).$$

Hence the set $\{\zeta^{\lambda_k}\}$ is complete in $A^2(\Delta)$ if and only if the set $\{\xi^{\lambda_k+1}e^{-\xi}\}$ is complete in $L^2((0, +\infty), d\xi/\xi)$, that is, the set $\{\xi^{\lambda_k+\frac{1}{2}}e^{-\xi/2}\}$ is complete in $L^2((0, +\infty))$.

Next, we consider the transformation $\xi \mapsto \xi^a = t$ of $(0, +\infty)$ onto itself and the induced isometry Λ of $L^2((0, +\infty))$ onto itself given by

$$(\Lambda\psi)(t) = \sqrt{\frac{1}{a}} \psi(t^{1/a}) \left(t^{\frac{1}{2}(\frac{1}{a}-1)}\right).$$

Under such a transformation, since $\lambda_k = ak + c_0 + ib$, we see that $\{\xi^{\lambda_k+\frac{1}{2}}e^{-\xi/2}\}$ is complete in $L^2((0, +\infty))$ if and only if $\{t^k t^\alpha e^{-\frac{1}{2}t^{1/a}}\}$ is complete in $L^2((0, +\infty))$, where

$$\alpha = \frac{c_0 + 1}{a} - \frac{1}{2} + i\frac{b}{2a}.$$

We know from [Sze75, Thm. 5.7.1] that the system $\{t^{n+c}e^{-t/2} : n = 0, 1, 2, \dots\}$, with $c > -1/2$, is complete in $L^2((0, +\infty))$. Thus, if $\psi \in L^2((0, +\infty))$ is orthogonal to $t^k t^\alpha e^{-\frac{1}{2}t^{1/a}}$ for all $k = 0, 1, 2, \dots$ it follows that

$$\int_0^{+\infty} t^{k+\operatorname{Re} \alpha} e^{-t/2} \overline{t^{-i\operatorname{Im} \alpha} e^{\frac{1}{2}(t-t^{1/a})}} \psi(t) dt = 0$$

for $k = 0, 1, 2, \dots$. Since $0 < a < 1$, $e^{\frac{1}{2}(t-t^{1/a})}$ is bounded and also $\operatorname{Re} \alpha > -\frac{1}{2}$, we obtain that $t^{-i\operatorname{Im} \alpha} e^{\frac{1}{2}(t-t^{1/a})} \psi = 0$, that is, $\psi = 0$. This concludes the proof. \square

3. COMPLETE SETS IN $A^2(\mathcal{W}'_\mu)$

From our Müntz–Szász Theorem 2.1 for the Bergman space $A^2(\Delta)$, we obtain the following density result in $A^2(\mathcal{W}'_\mu)$.

Theorem 3.1. *Let $\mu > \pi/2$. Let $H_{\ell,j}(z_1, z_2)$ be as in (13). Then $\{H_{\ell,j}\}_{\ell,j \in \mathbb{Z}, \ell \geq 0}$, is a complete set in $A^2(\mathcal{W}'_\mu)$.*

Notice that the set $\{H_{\ell,j}\}$, $\ell, j \in \mathbb{Z}$, $\ell \geq 0$, is the union of the two sets in (14).

Proof. We wish to show that if $F \in A^2(\mathcal{W}'_\mu)$ is orthogonal to $H_{\ell,j}$, for $\ell, j \in \mathbb{Z}$, $\ell \geq 0$, then F is identically zero. It suffices to show that, for each $j \in \mathbb{Z}$ fixed, any function $F \in \mathcal{H}^j$ orthogonal to $H_{\ell,j}$ for all $\ell \geq 0$, is identically zero.

Writing $F(z_1, z_2) = f(z_1, |z_2|)z_2^j$, from (11) we then have

$$\begin{aligned} 0 &= \langle F, H_{\ell,j} \rangle_{A^2(\mathcal{W}'_\mu)} = \pi \int_{\Delta} \int_{|s| < \mu} f(\zeta e^{is}, e^{s/2}) \overline{E_{c_0 + \nu\ell + i(j+1)/2}(\zeta e^{is}, e^{s/2})} e^{s(j+1)} ds dA(\zeta) \\ &= \pi \int_{\Delta} \int_{|s| < \mu} f(\zeta e^{is}, e^{s/2}) e^{s[(j+1)/2 + i(c_0 + \nu\ell)]} \overline{\zeta^{c_0 + \nu\ell + i(j+1)/2}} dA(\zeta), \end{aligned} \quad (15)$$

for $\ell = 0, 1, \dots$. Notice that the function

$$\begin{aligned} Tf(\zeta, w) &= \int_{|s| < \mu} f(\zeta e^{is}, e^{s/2}) e^{s[(j+1)/2 + i c_0]} e^{iw} ds \\ &= \mathcal{F}(f(\zeta e^{is}, e^{s/2}) e^{s[(j+1)/2 + i c_0]} \chi_{\{|s| < \mu\}})(w) \end{aligned} \quad (16)$$

is analytic in $\zeta \in \Delta$, and by the Paley–Wiener theorem [PW87], is an entire function in w of exponential type at most μ . Moreover, the function

$$w \mapsto \pi \int_{\Delta} Tf(\zeta, w) \overline{\zeta^{c_0 + \nu\ell + i(j+1)/2}} dA(\zeta)$$

is again an entire function of exponential type at most μ and by (15) it vanishes at the points $w_\ell = \nu\ell$. Observe that

$$\limsup_{r \rightarrow +\infty} \frac{\exp\{2 \sum_{\nu\ell < r} 1/(\nu\ell)\}}{r^{2\mu/\pi}} = \left(\frac{1}{\nu}\right)^{\frac{2}{\nu}} \lim_{r \rightarrow +\infty} r^{1/\nu} = +\infty. \quad (17)$$

By a classical result of Fuchs [Fuc46], we know that an entire function of type μ whose zero set $\{w_\ell = \nu\ell\}$ satisfies (17) must vanish identically, that is,

$$\int_{\Delta} Tf(\zeta, w) \overline{\zeta^{c_0 + \nu\ell + i(j+1)/2}} dA(\zeta) = 0,$$

for $\ell = 0, 1, \dots$. Since $\mu > \pi/2$ we have $\nu < 1$ and by Theorem 2.1 it now follows that $Tf(\cdot, w) = 0$, hence,

$$\int_{|s| < \mu} f(\zeta e^{is}, e^{s/2}) e^{s[(j+1)/2 + i c_0]} e^{isw} ds = 0, \quad (18)$$

for all $\zeta \in \Delta$ and $w \in \mathbb{C}$. This implies that f vanishes identically and we are done. \square

Notice that, had we considered either of the orthogonal systems mentioned in Corollary 1.5, we would have ended up with the points $\{w_{2\ell}\}$ only, or with $\{w_{2\ell+1}\}$. The analog of Condition (17)

would not have been satisfied and we could not have proved completeness using this approach. In fact, we are going to show in the next proposition that each of the two systems is incomplete.

It is also worth mentioning that the worm domains \mathcal{W}'_μ are increasingly badly behaved as μ becomes large. On the other hand, the proof of our density result breaks down when $\mu \leq \pi/2$. This is somewhat surprising, since when $\mu \leq \pi/2$ the fibers over z_1 are connected, the geometry of the domain is much simpler and in principle it should be easier to obtain such results on \mathcal{W}'_μ when $\mu \leq \pi/2$.

Proposition 3.2. *Let $\{H_{\ell,j}\}_{\ell,j \in \mathbb{Z}, \ell \geq 0}$ be as in Theorem 3.1. Then, for each m fixed,*

$$\|H_{2m+1,j}\|_{A^2(\mathcal{W}'_\mu)}^2 > \sum_{j'=-\infty}^{+\infty} \sum_{k=0}^{+\infty} \frac{1}{\|H_{2k,j'}\|_{A^2(\mathcal{W}'_\mu)}^2} |\langle H_{2m+1,j}, H_{2k,j'} \rangle|^2, \quad (19)$$

and, analogously, for each k fixed,

$$\|H_{2k,j}\|_{A^2(\mathcal{W}'_\mu)}^2 > \sum_{j'=-\infty}^{+\infty} \sum_{m=0}^{+\infty} \frac{1}{\|H_{2m+1,j'}\|_{A^2(\mathcal{W}'_\mu)}^2} |\langle H_{2k,j}, H_{2m+1,j'} \rangle|^2.$$

Hence, neither system $\{H_{2k,j}\}$ nor $\{H_{2m+1,j}\}$ is complete in $A^2(\mathcal{W}'_\mu)$.

Proof. By orthogonality, it suffices to consider the case $j' = j$ and, dropping the index j , we write $F_k = H_{2k,j}$ and $G_m = H_{2m+1,j}$. By Proposition 1.3 we have that

$$\begin{aligned} \langle G_m, F_k \rangle_{A^2(\mathcal{W}'_\mu)} &= (2\pi)^2 \frac{\sin[\mu(2k - (2m+1))\nu]}{(2k - (2m+1))\nu} \\ &\quad \times \frac{\Gamma(2c_0 + 2 + (2(k+m)+1)\nu)}{\Gamma(c_0 + 2 + 2k\nu + i\frac{j+1}{2})\Gamma(c_0 + 2 + (2m+1)\nu - i\frac{j+1}{2})}, \\ \|F_k\|_{A^2(\mathcal{W}'_\mu)}^2 &= (2\pi)^2 \mu \frac{\Gamma(2c_0 + 2 + 4k\nu)}{|\Gamma(c_0 + 2 + 2k\nu + i\frac{j+1}{2})|^2}, \\ \|G_m\|_{A^2(\mathcal{W}'_\mu)}^2 &= (2\pi)^2 \mu \frac{\Gamma(2c_0 + 2 + 2(2m+1)\nu)}{|\Gamma(c_0 + 2 + (2m+1)\nu - i\frac{j+1}{2})|^2}. \end{aligned}$$

Therefore, (19) is equivalent to

$$\begin{aligned} &\mu\Gamma(2c_0 + 2 + 2(2m+1)\nu) \\ &> \sum_{k=0}^{+\infty} \left[\frac{\sin[\mu(2k - (2m+1))\nu]}{(2k - (2m+1))\nu} \right]^2 \frac{\Gamma(2c_0 + 2 + (2(k+m)+1)\nu)^2}{\mu\Gamma(2c_0 + 2 + 4k\nu)}, \end{aligned} \quad (20)$$

which in turn is implied by,

$$\begin{aligned} 1 &> \sum_{k=0}^{+\infty} \frac{1}{[\mu(2k - (2m+1))\nu]^2} \frac{\Gamma(2c_0 + 2 + (2(k+m)+1)\nu)^2}{\Gamma(2c_0 + 2 + 2(2m+1)\nu)\Gamma(2c_0 + 2 + 4k\nu)} \\ &= \sum_{k=0}^{+\infty} \frac{1}{\pi^2(k - \frac{2m+1}{2})^2} \frac{\Gamma(2c_0 + 2 + (2(k+m)+1)\nu)^2}{\Gamma(2c_0 + 2 + 2(2m+1)\nu)\Gamma(2c_0 + 2 + 4k\nu)}. \end{aligned} \quad (21)$$

Now, on the one hand the right-hand side in (21) is less than or equal to

$$\sum_{k=0}^{+\infty} \frac{1}{\pi^2 \left(k - \frac{2m+1}{2}\right)^2},$$

since for all $x, y > 0$, $c \geq 0$,

$$\begin{aligned} \Gamma(c+x+y)^2 &= \left(\int_0^{+\infty} t^{x+y+c-1} e^{-t} dt \right)^2 \\ &\leq \left(\int_0^{+\infty} t^{2x+c-1} e^{-t} dt \right) \left(\int_0^{+\infty} t^{2y+c-1} e^{-t} dt \right) = \Gamma(c+2x)\Gamma(c+2y). \end{aligned}$$

On the other hand, we claim that

$$\pi^2 > \sum_{k=0}^{+\infty} \frac{1}{\left(k - \frac{2m+1}{2}\right)^2}.$$

Indeed, setting $h(w) = \pi \cot(\pi w)$ and $Q(w) = \left(w - \frac{2m+1}{2}\right)^2$, we have

$$\operatorname{Res}\left(\frac{h}{Q}, k\right) = \frac{1}{Q(k)} = \frac{1}{\left(k - \frac{2m+1}{2}\right)^2}$$

for all $k \in \mathbb{Z}$ and

$$\operatorname{Res}\left(\frac{h}{Q}, \frac{2m+1}{2}\right) = \lim_{w \rightarrow \frac{2m+1}{2}} \frac{\pi}{\sin(\pi w)} \frac{\cos(\pi w)}{w - \frac{2m+1}{2}} = -\pi^2.$$

The fact that

$$0 = \lim_{n \rightarrow +\infty} \int_{\partial D(0, n+\frac{1}{2})} \frac{h(w)}{Q(w)} dw = 2\pi i \lim_{n \rightarrow +\infty} \left[\operatorname{Res}\left(\frac{h}{Q}, \frac{2m+1}{2}\right) + \sum_{k=-n}^n \operatorname{Res}\left(\frac{h}{Q}, k\right) \right]$$

implies that

$$\pi^2 = \sum_{k=-\infty}^{+\infty} \frac{1}{\left(k - \frac{2m+1}{2}\right)^2} > \sum_{k=0}^{+\infty} \frac{1}{\left(k - \frac{2m+1}{2}\right)^2},$$

as claimed. This concludes the proof. \square

Finally, we show that the complete system of Theorem 3.1 is not a basis for $A^2(\mathcal{W}'_\mu)$, for all $\mu \geq \pi/2$.

Theorem 3.3. *Let $\mu \geq \pi/2$, and let $H_{\ell,j}(z_1, z_2) = E_{c_0+\nu\ell+i(j+1)/2}(z_1, z_2)z_2^j$, $\ell, j \in \mathbb{Z}$, $\ell \geq 0$. For each $j \in \mathbb{Z}$ fixed, the function $H_{0,j}$ is in the $A^2(\mathcal{W}'_\mu)$ -closure of $\operatorname{span}\{H_{\ell,j}, \ell = 1, 2, \dots\}$.*

Proof. We first assume that $\mu > \pi/2$.

Let $Q = Q_n$ be a polynomial of degree n of one complex variable, without constant term, $Q(w) = \sum_{\ell=1}^n c_\ell w^\ell$. Then, arguing as in (7) we have

$$\begin{aligned}
& \|H_{0,j} - \sum_{\ell=1}^n c_\ell H_{\ell,j}\|_{A^2(\mathcal{W}'_\mu)}^2 \\
&= \pi \int_{|s| < \mu} \int_{\Delta} \left| \zeta^{c_0+i(j+1)/2} e^{is(c_0+i(j+1)/2)} - \sum_{\ell=1}^n c_\ell \zeta^{c_0+\nu\ell+i(j+1)/2} e^{is(c_0+i(j+1)/2+\nu\ell)} \right|^2 dA(\zeta) e^{s(j+1)} ds \\
&= \pi \int_{|s| < \mu} \int_{\Delta} \left| \zeta^{c_0+i(j+1)/2} \right|^2 \left| 1 - \sum_{\ell=1}^n c_\ell \zeta^{\nu\ell} e^{is\nu\ell} \right|^2 dA(\zeta) ds \\
&\leq C \int_{|s| < \mu} \int_{\Delta} \left| 1 - \sum_{\ell=1}^n c_\ell (\zeta^\nu e^{is\nu})^\ell \right|^2 dA(\zeta) ds \\
&= C \frac{1}{\nu^2} \int_{|s| < \mu} \int_{\Omega_s} \left| 1 - \sum_{\ell=1}^n c_\ell w^\ell \right|^2 |w^{\frac{1}{\nu}-1}|^2 dA(w) ds, \tag{22}
\end{aligned}$$

where we have set $w = \zeta^\nu e^{is\nu}$. Since $\mu > \pi/2$, $\nu < 1 - 2\delta$ for some $\delta > 0$, so that $-\frac{\pi}{2} < \nu s < \frac{\pi}{2}$ and $0 < \nu\frac{\pi}{2} < \frac{\pi}{2}(1 - 2\delta)$. Hence,

$$\begin{aligned}
\Omega_s &\subseteq \{w = \rho e^{it} : 0 < \rho < 2^\nu, \nu(s - \frac{\pi}{2}) < t < \nu(s + \frac{\pi}{2})\} \\
&\subseteq \{w = \rho e^{it} : 0 < \rho < 2^\nu, |t| < \pi(1 - \delta)\} \\
&=: S.
\end{aligned}$$

Plugging this into (22) we obtain that

$$\begin{aligned}
\|H_{0,j} - \sum_{\ell=1}^n c_\ell H_{\ell,j}\|_{A^2(\mathcal{W}'_\mu)}^2 &\leq C \int_{|s| < \mu} \int_S |1 - Q_n(w)|^2 |w|^{2(\frac{1}{\nu}-1)} dA(w) ds \\
&= C \int_S |1 - Q_n(w)|^2 |w|^{2(\frac{1}{\nu}-1)} dA(w).
\end{aligned}$$

Setting $d\omega(w) = |w|^{2(\frac{1}{\nu}-1)} dA(w)$, the conclusion will follow if we show that there exist polynomials $P_n = 1 - Q_n$ such that $P_n(0) = 1$ and $\|P_n\|_{A^2(S, d\omega)} \rightarrow 0$ as $n \rightarrow +\infty$.

In order to prove that such polynomials exist, let Δ_+ be the half disk $\{z \in \mathbb{C} : |z| < 1, \operatorname{Re} z > 0\}$ and $p(z) = (z - \frac{1}{2})^2 + \frac{3}{4}$. Then $p(0) = 1$, and $|p(z)| \leq 1$ for $z \in \overline{\Delta_+}$, as it is elementary to check. Therefore, $F(w) = p((2^{-\nu}w)^{1/[2(1-\delta)]})$ is a function holomorphic on S such that

- F is continuous on \overline{S} ;
- $F(0) = 1$;
- $|F(w)| \leq 1$ on \overline{S} .

Observe that $\omega(S) < +\infty$. Given $\varepsilon > 0$, let K be a compact subset of S such that $\omega(S \setminus K) < \varepsilon$ and let n be a positive integer such that $|F^n(w)| \leq \varepsilon$ for $w \in K$. Then

$$\int_S |F^n(w)|^2 d\omega(w) \leq C\varepsilon.$$

By Mergelyan's approximation theorem (see [Rud66] e.g.), we can find polynomials p_n such that $|F^n(w) - p_n(w)| \leq \varepsilon$ for $w \in \overline{S}$. Finally, we set $P_n = \frac{1}{p_n(0)} p_n$ and the conclusion follows easily.

Finally, let $\mu = \pi/2$, so that $\nu = 1$. Set $\mathcal{D} = \cup_{|s| < \pi/2} \{z_1 : |z_1 - e^{is}| < 1\}$. We have,

$$\begin{aligned} \|H_{0,j} - \sum_{\ell=1}^n c_\ell H_{\ell,j}\|_{A^2(\mathcal{W}'_\mu)}^2 &= \pi \int_{|s| < \mu} \int_{|z_1 - e^{is}| < 1} |z_1^{2c_0 + i(j+1)}| \left| 1 - \sum_{\ell=1}^n c_\ell z_1^\ell \right|^2 dA(z_1) e^{s(j+1)} ds \\ &\leq C \int_{\mathcal{D}} \left| 1 - \sum_{\ell=1}^n c_\ell z_1^\ell \right|^2 dA(z_1). \end{aligned} \quad (23)$$

We observe that \mathcal{D} is a Jordan domain, having the origin as a boundary point. By [Far34] we know that the polynomials are dense in $A^2(\mathcal{D})$ and by [Bre85] it follows that there exists no bounded boundary evaluation point on the space of polynomials. Hence, the right hand side of (23) can be made arbitrarily small and the conclusion now follows. We leave the simple details to the reader. \square

CONCLUDING REMARKS

Thanks to work of several authors, the worm domain has become an important object of study. In particular, we are beginning to understand the Bergman kernel and projection on some versions of the worm. But the original smooth worm \mathcal{W}_μ is particularly resistive to analysis. It does not have the built-in symmetries of some of the non-smooth worms. In particular, we do not have a useful complete orthogonal basis for the Bergman space on \mathcal{W}_μ . In addition to the alternative approach mentioned in [KPS15, §5], this paper has offered some first steps towards addressing that problem.

REFERENCES

- [Bar92] David Barrett. Behavior of the Bergman projection on the Diederich-Fornaess worm. *Acta Math.*, 168(1-2):1–10, 1992.
- [BBG⁺04] David Békollé, Aline Bonami, Gustavo Garrigós, Cyrille Nana, Marco M. Peloso, and Fulvio Ricci. Lecture notes on Bergman projectors in tube domains over cones: an analytic and geometric viewpoint. *IMHOTEJ. Afr. Math. Pures Appl.*, 5:Exp. I, front matter + ii + 75, 2004.
- [Bre85] James E. Brennan. Weighted polynomial approximation, quasianalyticity and analytic continuation. *J. Reine Angew. Math.*, 357:23–50, 1985.
- [Cat80] David Catlin. Boundary behavior of holomorphic functions on pseudoconvex domains. *J. Differential Geom.*, 15(4):605–625 (1981), 1980.
- [Chr96] Michael Christ. Global C^∞ irregularity of the $\bar{\partial}$ -Neumann problem for worm domains. *J. Amer. Math. Soc.*, 9(4):1171–1185, 1996.
- [CS01] So-Chin Chen and Mei-Chi Shaw. *Partial Differential Equations in Several Complex Variables*, volume 19. American Mathematical Society, Providence, RI, 2001.
- [DF77] Klas Diederich and John Erik Fornaess. Pseudoconvex domains: an example with nontrivial Nebenhülle. *Math. Ann.*, 225(3):275–292, 1977.
- [DGGMR07] Peter Duren, Eva A. Gallardo-Gutiérrez, and Alfonso Montes-Rodríguez. A Paley-Wiener theorem for Bergman spaces with application to invariant subspaces. *Bull. Lond. Math. Soc.*, 39(3):459–466, 2007.
- [Far34] O. J. Farrell. On approximation to an analytic function by polynomials. *Bull. Amer. Math. Soc.*, 40(12):908–914, 1934.
- [Fuc46] W. H. J. Fuchs. A generalization of Carlson’s theorem. *J. London Math. Soc.*, 21:106–110, 1946.
- [GR96] I. S. Gradshteyn and I. M. Ryzhik. *Table of integrals, series, and products*. Academic Press Inc., San Diego, CA, fifth edition, 1996. CD-ROM version 1.0 for PC, MAC, and UNIX computers.

- [Kis91] Christer O. Kiselman. A study of the Bergman projection in certain Hartogs domains. In *Several complex variables and complex geometry, Part 3 (Santa Cruz, CA, 1989)*, volume 52 of *Proc. Sympos. Pure Math.*, pages 219–231. Amer. Math. Soc., Providence, RI, 1991.
- [KP08a] Steven G. Krantz and Marco M. Peloso. Analysis and geometry on worm domains. *J. Geom. Anal.*, 18(2):478–510, 2008.
- [KP08b] Steven G. Krantz and Marco M. Peloso. The Bergman kernel and projection on non-smooth worm domains. *Houston J. Math.*, 34(3):873–950, 2008.
- [KPS15] Steven G. Krantz, Marco M. Peloso, and Caterina Stoppato. Bergman kernel and projection on the unbounded Diederich–Fornæss worm domain. *Ann. Sc. Norm. Super. Pisa Cl. Sci.*, DOI: 10.2422/2036-2145.201503.012, 2015.
- [Mün14] C. H. Müntz. Über den Approximationssatz von Weierstrass. In *H. A. Schwarz’s Festschrift*, pages 303–312. Berlin, 1914.
- [PW87] Raymond E. A. C. Paley and Norbert Wiener. *Fourier transforms in the complex domain*, volume 19 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 1987. Reprint of the 1934 original.
- [Rud66] Walter Rudin. *Real and Complex Analysis*. McGraw-Hill, Inc., Publishers, New York, 1966.
- [Szá16] Otto Szász. Über die Approximation stetiger Funktionen durch lineare Aggregate von Potenzen. *Math. Ann.*, 77(4):482–496, 1916.
- [Sze75] Gábor Szegő. *Orthogonal polynomials*. American Mathematical Society, Providence, RI, fourth edition, 1975. American Mathematical Society, Colloquium Publications, Vol. XXIII.

CAMPUS BOX 1146, WASHINGTON UNIVERSITY IN ST. LOUIS, ST. LOUIS, MISSOURI 63130

DIPARTIMENTO DI MATEMATICA “F. ENRIQUES”, UNIVERSITÀ DEGLI STUDI DI MILANO, VIA C. SALDINI 50, I-20133 MILANO

ISTITUTO NAZIONALE DI ALTA MATEMATICA, UNITÀ DI RICERCA DI FIRENZE C/O DIMAI “U. DINI” UNIVERSITÀ DI FIRENZE, VIALE MORGAGNI 67/A, I-50134 FIRENZE

E-mail address: `sk@math.wustl.edu`

E-mail address: `marco.peloso@unimi.it`

E-mail address: `stoppato@math.unifi.it`